



OPEN

Different organic manure sources and NPK fertilizer on soil chemical properties, growth, yield and quality of okra

Aruna Olasekan Adekiya¹✉, Wutem Sunny Ejue¹, Adeniyi Olayanju², Oluwagbenga Dunsin¹, Christopher Muiyiwa Aboyeji¹, Charity Aremu¹, Kehinde Adegbite¹ & Olanike Akinpelu¹

Use of organic manures to meet the nutrient requirement of crop would be an inevitable practice in the years to come for sustainable agriculture since organic manures generally improve the soil physical, chemical and biological properties. Hence, field experiments were carried out in 2017 and 2018 to compare the impact of different organic manures and NPK fertilizer on soil properties, growth, yield, proximate and mineral contents of okra (*Abelmoschus esculentus* L.). The treatments each year 2017 and 2018 consisted of: rabbit manure, cow dung, poultry manure, green manure [Mexican sunflower (*Tithonia diversifolia* Asteraceae)], pig manure, NPK 15-15-15 fertilizer applied at 120 kg N ha⁻¹ and a control (no manure/inorganic fertilizer). The seven treatments were laid out in a randomized complete block design with three replication. Organic manures and NPK fertilizer increased the soil organic matter (OM), N, P, K, Ca and Mg (NPK fertilizer did not increase OM, Ca and Mg significantly), growth, yield, minerals, protein, ash, carbohydrate and mucilage contents of okra fruit as compared with control. Organic manures improved okra yield compared with NPK fertilizer. Okra growth and yield parameters were significantly higher in 2018 compared with 2017. Control, rabbit manure, cow dung, poultry manure, green manure, pig manure and NPK fertilizer in 2018 increased the pod yield of okra by 9.7%, 35.3%, 57.9%, 36.2%, 39.2%, 45.5% and 3.2%, respectively compare with the same treatment in 2017. Amongst various organic manures, poultry manure produced significantly higher plant growth, yield, mineral and proximate composition of okra because of its high soil chemical properties which could be related to its lowest C: N ratio, lignin and lignin: N ratio. Results also showed that okra grown during high intensity rainfall has higher yield but with reduced quality except its mucilage content. Therefore, planting of okra with poultry manure under moderate rainfall will enhance the health benefit from the fruit, however, those that desire its mucilage content planting during high rainfall is recommended.

Okra (*Abelmoschus esculentus* L.) is an important tropical and subtropical vegetable crop grown for its fresh leaves, buds, flowers, pods, stems and seeds. The fresh pods can be eaten as vegetables in form of salads, in soups and stews or boiled. Okra fruits contain mucilage upon cooking. The mucilage has medical importance as it is used as a plasma replacement or blood volume expander and binds cholesterol and bile acid carrying toxins dumped into it by the liver¹. Okra fruits contain fiber, vitamin C, folate and antioxidants². The seeds contain oil that is edible by man as well as in soap industry³.

Despite the enormous potentials of okra fruits production, its yield per hectare and quality in Nigeria had been greatly hampered by the low fertility status and organic matter contents of the soils which translate into low productivity and consequently reducing income for the farmer⁴. According to Adekiya et al.⁴, the yield of okra in Nigeria is currently very low about 2.7 t ha⁻¹ owing to low native soil fertility status among other factors. Lack of sufficient amounts of nutrients result in poor performance of the crop with growth been affected resulting to low yield. It has been reported that the maintenance of soil organic matter (OM) is the basis of sustainable crop

¹College of Agricultural Sciences, Landmark University, PMB 1001, Omu-Aran, Kwara State, Nigeria. ²Department of Agricultural & Biosystem Engineering, Landmark University, PMB 1001, Omu-Aran, Kwara State, Nigeria. ✉email: adekiya2009@yahoo.com

	2017	2018
Soil properties		
pH (H ₂ O)	6.10	6.20
OM (%)	2.56	2.44
Total N (%)	0.10	0.15
Available P (mg/kg)	9.50	8.90
Exchangeable cations (cmol kg⁻¹)		
K	0.15	0.14
Ca	1.80	1.90
Mg	0.41	0.38
Particle size distribution (%)		
Sand	76	74
Silt	13	15
Clay	11	11
Textural class	Sandy loam	Sandy loam

Table 1. Pre-planting physico-chemical characteristics of the soil (0–15 cm) at the experimental sites in 2017 and 2018.

production in Nigeria and other tropical countries⁵. Hence, there is need to improve the fertility of the soil for continuous and increased crop production.

In years to come, utilization of organic manure to meet crop nutrient requirement will be an unavoidable practice to enhance sustainable agriculture, this is because, the physical, chemical and biological properties of soil is generally improved by the addition of organic manures which in turn enhances crop productivity and maintains the quality of crop produce⁶. Although, in comparison to inorganic fertilizers, organic manures contain smaller quantities of plant nutrients. The use of inorganic fertilizer to increase yield has been found to be effective as a short-term solution but demands consistent use on a long-term basis. The high cost of inorganic fertilizers makes it uneconomical and out of reach to poor farmers and it is also undesirable due to its hazardous environmental effects⁷. Therefore, it is essential to investigate the use of locally sourced organic materials which are environment friendly, cheap and probably an effective way of improving and sustaining the productivity of soils and arable crops such as okra.

Okra yield responses to organic and inorganic fertilizers have been reported by several workers^{2,4,8–10}. However, since organic and inorganic fertilizers contain different chemical composition and quality they therefore may react differently when applied to the soil with respect to soil chemical properties, crop yield and quality. This aspect needs investigation especially in Nigeria where such data on the effects of different organic and inorganic amendments on the mineral and proximate contents are lacking. Available work dealt with organic amendment on the yield of okra without putting into consideration the quality of okra fruits. In Sokoto northern Nigeria¹⁰, investigated different sources of organic manure (cow, sheep and poultry manure) on growth and yield of okra. Their results revealed that poultry manure promotes higher growth and yield of okra compared with cow and sheep manure.

Fagwalawa and Yahaya¹¹ investigated the effects of Sheep, cow and poultry manures and their combinations on the growth and yield of okra, their result also revealed that poultry manure has the highest yield. Also in Malaysia⁹, six different treatments (no fertilizer, NPK fertilizer, poultry manure, rat manure, goat manure and rabbit manure) were investigated on the growth and yield of okra, According to the study, application of poultry manure significantly increased the growth and yields performances on okra compared to other types of organic fertilizers. To have a holistic approach on the response of organic manures on soil chemical properties and okra performance, data on the quality of such okra fruit is imperative. Therefore, the study is to compare the impact of different organic manures and NPK fertilizer on soil properties, growth, yield, proximate and mineral contents of okra grown in derived savanna zone of Nigeria.

Results

Pre-planting chemical and physical analysis of the experimental field and chemical composition of organic manures used for the experiment. The result of the pre-planting chemical and physical analysis of the experimental field (Table 1), indicated in both years that the soil was sandy loam and slightly acidic with a pH of 6.10 and 6.20 respectively in 2017 and 2018. The soil organic matter, total N available P and exchangeable Ca and K were low with both a little below the critical levels of 3.0%, 0.20%, 10.0 mg kg⁻¹, 2.0 cmol kg⁻¹ and 0.15 cmol kg⁻¹ respectively in both cropping seasons of 2017 and 2018¹². Exchangeable Mg was adequate in 2017 but low in 2018. Analysis of the soil amendments used for this experiment is shown in Table 2 with poultry manure having the highest percentage of N, P, K, Ca, Mg and the lowest lignin, lignin: N ratio, C: N ratio and organic C.

Response of soil chemical properties and bulk density to different organic manure sources and NPK fertilizer. Table 3 shows the result of the effect of different organic amendments and NPK fertilizer

Manure	Organic C (%)	N (%)	C: N	Lignin (%)	Lignin: N	P (%)	K (%)	Ca (%)	Mg (%)
Rabbit manure	30.1a	1.01e	29.8a	14.8a	14.7a	0.54b	1.95d	1.15d	0.40c
Cow dung	26.5b	1.86d	14.24b	14.6a	7.8b	0.82a	2.11c	1.01d	0.51b
Poultry manure	17.8e	2.91a	6.12e	6.1c	2.1e	0.84a	3.79a	3.34a	0.64a
Green manure	23.6c	2.51b	9.40d	8.1b	3.2d	0.52b	3.04b	3.01b	0.10d
Pig manure	20.1d	2.16c	9.77c	7.9b	3.7c	0.80a	2.16c	1.45c	0.54b

Table 2. Analysis of organic manures. Values followed by similar letters under the same column are not significantly different at $p=0.05$ according to Duncan's multiple range test.

Year	Amendments	pH (water)	OM (%)	N (%)	P (mg kg ⁻¹)	K (cmol kg ⁻¹)	Ca (cmol kg ⁻¹)	Mg (cmol kg ⁻¹)	Bulk density (g cm ⁻³)	
2017	Control	6.02a	2.47f	0.09f	7.9f	0.14g	2.81e	0.38g	1.55a	
	Rabbit	5.84ab	3.87a	0.14d	9.6e	0.19e	3.28d	0.40ef	1.20b	
	Cow dung	5.81ab	3.57bc	0.19c	10.3d	0.20de	4.80c	0.46d	1.19b	
	Poultry	5.85ab	3.08e	0.25a	27.4a	0.70a	5.84a	0.81a	1.19b	
	Green manure	5.87ab	3.23d	0.24a	15.9b	0.50b	4.96b	0.56b	1.18b	
	Pig manure	5.82ab	3.47c	0.21b	12.3c	0.26c	4.84c	0.49cd	1.18b	
	NPK	5.51b	2.50f	0.13e	9.5e	0.16f	2.91e	0.39fg	1.54a	
2018	Control	6.01a	2.39e	0.08f	7.2g	0.12f	2.69f	0.36e	1.55a	
	Rabbit	5.81ab	3.41a	0.17d	10.6e	0.22d	3.66d	0.45d	1.19b	
	Cow dung	5.81ab	3.11b	0.21c	13.4d	0.22d	5.15c	0.51c	1.19b	
	Poultry	5.79ab	2.95d	0.27a	29.1a	0.75a	6.96a	0.88a	1.19b	
	Green manure	5.77ab	3.07c	0.27a	19.4b	0.55b	5.48b	0.61b	1.19b	
	Pig manure	5.0ab	3.09c	0.24b	15.4c	0.31c	5.10c	0.53c	1.18b	
	NPK	5.51b	2.41e	0.13e	9.3f	0.15e	2.72e	0.37e	1.54a	
	Year (Y)	Ns	*	*	*	*	*	*	*	Ns
	Manure (M)	Ns	*	*	*	*	*	*	*	*
Y × M	Ns	*	*	*	*	*	*	*	Ns	

Table 3. Post-Harvest soil chemical properties. Values followed by similar letters under the same column are not significantly different at $p=0.05$ according to Duncan's multiple range test. *Significant difference at $p=0.05$; ns, not significant at 0.05.

on soil chemical properties. In both years, all organic sources of soil amendment increased soil OM, N, P, K, Ca and Mg significantly ($p < 0.05$) with respect to the control. NPK fertilizer did not increase OM, Ca and Mg contents of the soil relative to the control but increased N, P and K. Among organic manure sources, poultry manure has the highest values of all soil nutrients (except SOM) which was closely followed by green manure treatment with rabbit manure having the least values. The order of SOM among organic manures was: Rabbit manure > cow dung > Pig manure > green manure > Poultry manure. All organic manure sources increased soil N, P and K compared with NPK fertilizer. In both years, the control has the highest value of pH while NPK fertilizer has the least value, there were no significant differences between the pH values of control, rabbit manure, cow dung, poultry manure, green manure and pig manure. Also there were no significant differences in the pH values between organic manures and NPK fertilizer, but there was a significant difference between the pH values of the control and NPK fertilizer. Soil chemical properties in year 2018 were increased compared with year 2017 except soil pH. Soil bulk density in both years was significantly reduced in organic manure soils compared with the control and NPK fertilizer which has similar values. All organic manures have statistically similar values of bulk density. The interactions between year and amendment (Y × M) were significant for all soil properties except soil pH and bulk density.

Response of okra growth and yield to different organic manure sources and NPK fertilizer. The results of the response of okra growth and yield to different organic manure sources and NPK fertilizer are presented in Table 4. In 2017 and 2018, all amendments (both organic and inorganic) increased the growth (plant height and leaf area) and yield (okra pod weight and number of pods per plant) of okra compared with the control. Also in both years, poultry manure has the highest values followed by green manure. The order of okra pod yield was: poultry manure > green manure > pig manure > cow dung > NPK fertilizer = rabbit manure > control. Okra growth and yield parameters were significantly higher in 2018 compared with 2017. Control, rabbit manure, cow dung, poultry manure, green manure, pig manure and NPK fertilizer in 2018 increased the pod yield of okra by 9.7%, 35.3%, 57.9%, 36.2%, 39.2%, 45.5% and 3.2%, respectively compare with the same treat-

Year	Amendments	Yield (t ha ⁻¹)	No. of pods/plant	Plant height (m)	Leaf area (m ²)
2017	Control	3.1f	10g	0.29f	0.46e
	Rabbit	3.4e	12f	0.37e	0.54d
	Cow dung	3.8d	14de	0.41d	0.57d
	Poultry	5.8a	23a	0.56a	0.81a
	Green manure	5.1b	21b	0.51b	0.71b
	Pig manure	4.4c	17c	0.47c	0.64c
	NPK	3.5e	13ef	0.38e	0.55d
2018	Control	3.4g	12h	0.30f	0.50f
	Rabbit	4.6e	15e	0.41de	0.60e
	Cow dung	5.2d	16de	0.44d	0.65de
	Poultry	7.9a	26a	0.62a	0.94a
	Green manure	7.1b	24b	0.55b	0.80b
	Pig manure	6.4c	18c	0.50c	0.71c
	NPK	3.6f	14fg	0.39e	0.60e
	Year (Y)	*	*	*	*
	Manure (M)	*	*	*	*
	Y × M	*	*	*	*

Table 4. Growth and yield parameters. Values followed by similar letters under the same column are not significantly different at $p=0.05$ according to Duncan's multiple range test. *Significant difference at $p=0.05$.

Year	Amendment	Protein (%)	Ash (%)	Fat (%)	Carbohydrate (%)	Mucilage (%)	N (%)	P (%)	K (%)	Ca (%)	Mg (%)
2017	Control	9.91g	5.6f	12.9a	22.0d	6.35g	0.82e	10.6d	0.88e	2.41e	2.10e
	Rabbit	19.1e	6.2e	13.1ab	26.8c	7.34de	2.87d	12.3b	1.10c	3.22c	3.22c
	Cow dung	19.3de	6.4de	13.9ab	27.0c	7.58c	2.96d	12.6b	1.06c	3.10c	3.32c
	Poultry manure	23.5a	7.1a	10.4c	32.0a	9.88a	4.41a	13.4a	1.87a	4.43a	4.31a
	Green manure	22.1b	6.9ab	10.9c	30.1ab	9.57ab	4.21ab	12.9a	1.44b	3.61b	4.01b
	Pig manure	19.6c	6.1e	10.2c	29.0b	7.03e	3.11c	11.6c	1.74ab	3.55b	3.10c
	NPK	17.5f	6.7cd	11.9b	27.1c	6.89f	3.21c	11.4c	0.94d	2.96d	2.22d
2018	Control	9.69c	5.1f	12.10a	19.9e	6.84d	0.75d	9.31c	0.81c	2.51e	2.00c
	Rabbit	18.6b	5.6e	11.9ab	23.3d	7.58bc	2.65c	10.0b	0.96b	2.76c	2.80b
	Cow dung	18.91b	5.8de	11.6ab	24.2cd	7.99b	2.65c	10.9b	0.90b	2.75c	2.88b
	Poultry manure	20.88a	6.6a	9.6c	28.0a	10.7a	3.24a	11.9a	1.16a	3.26a	3.68a
	Green manure	20.55a	6.5ab	9.8c	27.6ab	10.4a	3.04a	11.8a	0.99b	2.91b	3.61a
	Pig manure	18.51ab	5.9d	9.2c	26.2b	7.8b	2.85b	10.6b	1.10ab	2.96b	2.86b
	NPK	17.20b	6.0cd	11.0b	23.0d	7.1c	2.86b	10.1b	0.99b	2.61de	2.21c
	Year (Y)	*	*	*	*	*	*	*	*	*	*
	Manure (M)	*	*	*	*	*	*	*	*	*	*
Y × M	*	*	*	*	*	*	*	*	*	*	

Table 5. Proximate and mineral contents of okra. Values followed by similar letters under the same column are not significantly different at $p=0.05$ according to Duncan's multiple range test. *Significant difference at $p=0.05$.

ment in 2018. The interactions between year and amendment (Y × M) were significant for all growth and yield parameters.

Response of okra quality to different organic manure sources and NPK fertilizer. The effect of different organic manure sources on proximate and mineral content of okra in 2017 and 2018 are shown in Table 5. In 2017 and 2018 different organic manure sources and NPK fertilizer increased protein, ash, carbohydrates, mucilage N, P, K Ca and Mg contents of okra fruits compared with the control. Fat contents of okra fruits were reduced with different organic sources and NPK fertilizer compared with the control. Among different amendment applied, poultry manure had the highest values of protein, carbohydrate, mucilage, N, P, K, Ca and Mg contents. Green manure had statistically similar value with poultry manure in protein, ash, fat, carbohydrate, mucilage, N, P and Mg contents in 2017 and in 2018, ash, fat, carbohydrate, mucilage, N and P were statisti-

cally similar. Poultry manure and green manure increased protein, carbohydrate, mucilage, N, P, K, Ca, Mg and reduced fat compared with NPK fertilizer. 2018 cropping of okra reduced protein, ash, fat, carbohydrate, N, P, K, Ca, Mg and increased mucilage contents in the fruits compared with 2017. The interaction ($Y \times M$) was also significant for proximate and mineral contents of the okra fruit.

Discussion

The different organic manures increased the soil organic matter (SOM), N, P, K, Ca and Mg contents of the soil compared with control. This result is consistent with the analysis recorded for the amendments in the present study that they contain these nutrients and also attested to the fact that the soil was deficient in these nutrients (Table 1). These nutrients were released into the soil as the organic manures were decomposed. Studies by^{13,14} have shown that animal manures and green manure increased soil OM, N, P and CEC and this was attributed to the availability and adequate supply of organic matter. The slightly lower pH of organic amended soil compared with the control could be due to the fact that during microbial decomposition of the incorporated manures, organic acids may have been released, which neutralized the alkalinity of the manures, thereby lowering the pH of the soil below their initial values. Adekiya et al.⁴ observed a similar trend in their work on organic amendments of soils. NPK fertilizer has the lowest pH as a result of leaching of bases from soil surface. Poultry manure significantly produced the highest soil chemical properties, this could be related to its lowest C:N ratio, lignin and lignin: N ratio (Table 2) which favours quick mineralization and release of nutrients to the soil compared with other soil amendments. Consequently, the quality of an organic amendment is defined in term of the relative content of nutrients (especially nitrogen), lignin, lignin: N and the C:N ratio¹⁴. Rabbit manure increased SOM compared with other organic manures. The increase in SOM of rabbit manure could be related to its high C:N and lignin content. Plant constituents, such as lignin retard decomposition. Organic materials with high C:N and lignin generally would favor nutrient immobilization, organic matter accumulation and humus formation¹⁴.

In both years, poultry manure increased N, P, K, Ca and Mg compared with other manures. This was due to the fact that poultry manure was low in C:N ratio, lignin and lignin/N and high in N, P, K, Ca and Mg compared with other manures. Due to high quality of this amendment, it decomposes quickly and release nutrients to the soil. The significant increase in soil nutrients especially OM, N, P and K in organic amendments compared with NPK fertilizer was due to leaching in NPK fertilizer treated plots. The reduction in Ca and Mg in NPK fertilizer plots compared with organically amended plot was due to the fact that NPK fertilizer did not contain Ca and Mg. The reduction of soil bulk density observed in both years with organic manures compared with control and NPK fertilizer could be attributed to increase in soil organic matter resulted from the degraded organic residues by soil microorganisms. Organic matter is known to improve soil structure, aeration and reduce soil bulk density¹⁵.

The decreasing order of okra yield were poultry manure > green manure > pig manure > cow dung > NPK fertilizer = rabbit manure > control. The positive effect of organic manure on growth and okra yield could be due to the contribution made by amendments to fertility status of the soils as the soils were low in organic carbon content. Manure when decomposed increases both macro and micro nutrients as well as enhances the physico-chemical properties of the soil for the betterment of okra growth.

Okra grown on poultry manure performed better in terms of growth and yield compared with other sources of organic soil amendment and NPK fertilizer. This could also be related to low C:N ratio, lignin and lignin/N values. These attributes of poultry manure will lead to fast mineralization and early release of nutrients to a short gestation crop like okra, hence there was a boost in the morphological growth of the plant which translate to greater yield compared with other amendments. Wolf and Snyder¹⁶ reported that C:N ratio of organic materials markedly influences the decomposition rate and the mineralization of N because N determines the growth and turnover of the microorganisms that mineralize organic carbon.

The reduced growth and yield of okra in plots treated with other manures in comparison with poultry manure could be as a result of immobilization of soil nutrients which occurs when soils are treated with animal manure of high lignin contents which results from the feed eaten by the animal¹⁷.

Higher yields were observed in poultry manure plots compared with inorganic fertilizer because, the nitrogen content of poultry manure is released to the soil gradually and steadily over longer time for the growth of the plant compared with nitrogen from NPK fertilizer which is prone to losses by run-off, volatilization, leaching and/or denitrification. Poultry manure has been said to be a better soil amendment compared with chemical fertilizers because of the greater capability of poultry manure to preserve its N¹⁸. The superior N supply by poultry manure during okra cropping in this experiment may be the reason for better growth and yield of okra in plots with poultry manure¹⁸. The obtained results corroborated the finding of¹⁹ that poultry manure increased the height of okra relative to other amendments.

The better performance of okra under NPK fertilizer plots compared with the control was due to release of nutrients (N, P and K) from the fertilizer which are absorbed by the okra plants. Okra growth and yield in second year (2018) was better than that of first year (2017). This was due to the differences in the amount of rainfall between the two years (Table 6). Year 2017 had 1238 mm of rainfall while it was 1428 mm in 2018. There was higher moisture in the first few weeks (month of May) of incorporation of the manures in 2018 compared with 2017 which may have led to better and quicker decomposition of the organic materials in 2018. Soil biological activities which causes degradation of organic materials is severely limited during limited moisture, but with the onset of the rains (2018), there is a flush in microbial activity²⁰.

The fact that organic manures and NPK fertilizer increased okra mineral contents compared with the control was attributed to increased availability of the nutrients in soil as a result of the mineralization of the manures leading to increased uptake by okra plants. In both years, the correlation coefficient between soil and okra fruit N, soil and okra fruit P, soil and okra fruit K, and okra fruit Ca, soil and okra fruit Mg were all significant with R values of 0.83, 0.71, 0.66 0.81 and 0.88, respectively at $P < 0.05$. Poultry manure had the highest values of N, P,

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
2017												
Rainfall (mm)	0	0	97.28	92.46	158.24	174.49	71.11	179.33	231.39	119.88	0	114.04
Relative humidity (%)	54.1	44.9	69.7	81.0	85.3	89.9	91.9	93.4	92.3	85.2	71.3	56.9
Mean temp (°C)	29.1	31.2	31.9	29.9	28.9	27.9	26.9	26.3	26.5	27.8	29.9	28.4
2018												
Rainfall (mm)	0	12.19	157.73	96.26	214.88	228.6	160.02	95.25	248.66	195.07	19.55	0
Relative humidity (%)	34.3	63.7	78.8	85.2	87.5	90.2	92.7	92.7	91.3	88.4	76.4	47.1
Mean Temp (°C)	26.6	29.6	29.3	29.2	28.1	27.7	26.4	26.2	26.4	27.5	28.8	28.9

Table 6. Meteorological data of the study area. *Source* Meteorological unit, Teaching and Research Farm, Landmark University, Omu-Aran, Kwara State, Nigeria.

K, Ca and Mg in the okra fruit compared with other amendment and NPK fertilizer. This result is also consistent with the soil chemical properties and growth and yield for this treatment. The poultry manure may have improved the availability of nutrients to the crop by enhancing the mineralization and supply of readily available nutrients to the soil²¹.

The increased nitrogen to the soil by the incorporation of organic manures increased the nitrogen uptake by the okra fruit thereby increasing protein. This explains the reasons for increase in crude protein values between organic amendment and control and NPK fertilizer treatment. Poultry manure has the highest value of crude protein, this is also consistent with its high soil N level (Table 3). Nitrogen is a major constituent of chlorophyll, protein, amino acids, various enzymes, nucleic acids and many other compounds in the cell of plants²². The crude protein contents in all the organically amended soils were above the critical 13–17%^{23,24}. Therefore the organic materials sustained good nutritive quality of okra as opposed to NPK and control. The ash content in poultry manure treatment was significantly higher possibly because of the balanced nutrient in the manure, unlike NPK with inferior contents of N, P, and K and the control with lower concentration of nutrients. In this study, poultry manure gave the highest N, P and K concentrations of 3.04%, 11.9% and 1.16% in 2017 and 4.41% 13.4% and 1.87% in 2018 (Table 6) respectively. The fat content of poultry manure was lowered compared with others, this was due to its high protein content of poultry manure. It has also been reported²⁵ that there is a negative correlation between fat and protein content. High nitrogen application reduced fat and increased protein content. Organic amendments and NPK fertilizer increased the mucilage contents of okra compared with the control. This could be as a result of increase in NPK from these amendments. This could be attributed to the increase in D-galactose, L-rhamnose and D-galacturonic acid contents in okra fruits by the application of nutrients through organic and inorganic sources which might have resulted in increase of mucilage content²⁶. The mucilaginous polysaccharide in the okra is rich in uronic acid (65%) and consists of rhamnose, galactose, glucose, galacturonic acid and glucuronic acid in addition to 3.7% acetyl groups²⁷. Ahmad et al.²⁷ also reported that compost and NPK fertilizer application increased the mucilage of borage plant. The highest value of mucilage in poultry manure could be related to increased soil nutrients compared with other treatments. Okra produced under organic amendments has better qualities (N, P, K, Ca, Mg, protein, ash and mucilage) compared with NPK fertilizer and control. This is because organic manures not only increase soil nutrients but also improves the physical by prevention of erosion and leaching of nutrients (in this experiment reduce bulk density) and biological properties²⁸. Organic manures also contains both micro and macro nutrients unlike NPK fertilizer that contains only N, P and K. Lumpkin²⁹ also, was of the opinion that organically produced vegetables were of higher qualities than those produced using conventional methods. Cropping of okra in 2018 reduced protein, ash, fat, carbohydrate, N, P, K, Ca and Mg contents of okra fruit compared with 2017 cropping. This could be adduced to high rainfall in 2018 compared with 2017. High rainfall has been reported to reduce the nutrient and proximate content of vegetables³⁰. Climatic conditions have a great effect on the concentration of mineral in plants. Variation in temperature and rainfall have been reported to influence the chemical composition in plants^{30,31}.

Conclusion

Results of this experiment showed that organic manures and NPK fertilizer increased the soil chemical properties (NPK fertilizer did not increase OM, Ca and Mg significantly), growth, yield, minerals, protein, ash, carbohydrate and mucilage contents of okra fruit as compared with control. Organic manures improved okra yield compared with NPK fertilizer. Amongst various organic manures, poultry manure produced significantly higher plant growth, yield, mineral and proximate composition of okra because of its high soil chemical properties which could be related to its lowest C:N ratio, lignin and lignin: N ratio. Results also showed that okra grown during high intensity rainfall has higher yield but with reduced quality except its mucilage content. Therefore, planting of okra with poultry manure under moderate rainfall will enhance the health benefit from the fruit, however, those that desire its mucilage content planting during high rainfall is recommended.

Material and methods

Site description, treatments and experimental layout. During 2017 and 2018 cropping seasons, okra was grown in a field experiment at the Landmark University Teaching and Research Farm, Omu-Aran, Kwara state (8° 9' N, 5° 61' E), with altitude of 562 m above sea level. The rainfall pattern was bimodal with peaks

in August (Table 6). The total annual rainfall in the area was about 1238 mm in 2017 with mean air temperature of 28.7 °C and mean relative humidity of 83.9%. In 2018, the total annual rainfall in the area was about 1428 mm with mean air temperature of 30.0 °C and mean relative humidity of 77.4%. The soil at the site of the experiment is an Alfisol classified as Oxic Hap-lustalf or Luvisol². The experimental site falls under the derived savanna agro-ecological zone of North-Central Nigeria. Weeds in the experimental soil before cultivation included Mexican sunflower (*Tithonia diversifolia* Asteraceae) and Guinea grass (*Panicum maximum* Jacq). Different sites were used for the experiment in 2017 and 2018, but the two sites were very close to each other.

The treatments each year 2017 and 2018 consisted of: rabbit manure, cow dung, poultry manure, green manure [Mexican sunflower (*Tithonia diversifolia* Asteraceae)], pig manure, NPK 15-15-15 fertilizer and a control (no manure/ inorganic fertilizer). Both the organic manures and NPK fertilizer were applied at the rate of 120 kg N ha⁻¹³². This was equivalent to 800 kg ha⁻¹ for NPK 15-15-15, 11.9 t ha⁻¹ for rabbit manure, 5.5 t ha⁻¹ for cow dung, 4.1 t ha⁻¹ for poultry, 4.8 t ha⁻¹ for green manure and 5.6 t ha⁻¹ for pig manure. The seven treatments were laid out in a randomized complete block design (RCBD) with three replications. Each block consisted of 7 plots measuring 3 × 2 m with 1 m block and 0.5 m between plots.

Land preparation, incorporation of manures, sowing of okra seeds and application of NPK fertilizer. The experimental field was cleared manually using cutlass and thrashed removed from the site before mechanical (ploughing and harrowing) land preparation after which plots were marked out to the required plot size of 3 × 2 m. A hand-held hoe was used to incorporate the manures into the soil to a depth of approximately 20 cm. All animal manures were obtained from the livestock section of Landmark University Teaching and Research Farm. Fresh top Mexican sunflower was collected from a nearby farm and hedge containing green tender stems and the leaves. The plots were allowed for 4 weeks before sowing of okra seeds. Sowing of okra variety NHAe-47-4 was done late April for both 2017 and 2018 croppings. Two seeds of okra were sown per hole at inter-row spacing of 0.6 m and 0.6 m intra-row spacing manually giving a plant population of 27,778 plants ha⁻¹. At 2 weeks after sowing, thinning to one plant per stand was done and this was followed by manual weeding using hand hoe before treatment application. Subsequent weeding was done as needed. NPK 15-15-15 fertilizer was applied by side placement at about 8–10 cm away from the sown seeds two weeks after sowing. Insect pests were controlled by spraying cypermethrin weekly at the rate of 30 ml per 10 l of water from 2 weeks after sowing till 4 weeks after sowing.

Determination of soil properties. Before the start of the experiment, soil samples from topsoil (0–15 cm) were taken from random spots in the study area and were bulked together, air-dried and sieved using a 2-mm sieve and their physical and chemical characteristics were determined². Textural class of the soils were determined by the method of³³. Soil organic carbon (OC) was determined by the procedure of Walkley and Black using the dichromate wet oxidation method³⁴. Total N was determined by the micro-Kjeldahl digestion method³⁵. Available P was determined by Bray-1 extraction followed by molybdenum blue colorimetry³⁶. Exchangeable K, Ca, and Mg were extracted using 1 M ammonium acetate³⁷. Thereafter, concentration of K was determined on a flame photometer, and Ca and Mg were determined by EDTA titration method. Soil pH was determined using a soil–water medium at a ratio of 1:2 with a digital electronic pH meter. At the termination of the experiment in 2017 and 2018, soil samples were also collected (on plot basis) and similarly analysed for soil chemical properties as described above. Also at the end of the experiment each year, soil bulk density were determined all plots. Five undisturbed samples (0.04 m diameter, 0.15 m high) were collected at 0–0.15 m depth from the centre of each plot at random and 0.15 m away from each okra plant using core steel sampler⁴. The samples were used to evaluate bulk density after oven-drying at 100 °C for 24 h⁴.

Determination of growth and yield parameters. Collection of data for growth (plant height and leaf area) was done at mid-flowering of okra plant (about 47 days after sowing). Plant height was determined by the use of meter rule while leaf area was determined by using the model {LA = 0.34(LW)^{1.12}} developed by³⁸, where LA = leaf area, L = leaf length and W = leaf width. Edible pods were harvested at four day intervals, counted and weighed. Pod weight was evaluated based on the cumulative harvests per plot. For 2017 cropping of okra, fruits were harvested until November 2017 while for 2018 okra cropping, fruits were harvested until July 2018.

Chemical analysis of okra fruits and organic manures. At harvest in 2017 and 2018, eight okra fruits of uniform sizes were randomly collected from each plot and analyzed for mineral contents according to methods recommended by the Association of Official Analytical Chemists³⁹. One gram of each sample was digested using 12 cm⁻³ of the mix of HNO₃, H₂SO₄, and HClO₄ (7:2:1 v/v/v). Contents of N, P, K, Ca and Mg were determined by atomic absorption spectrophotometry. Lignin content of the organic amendments was determined from acid-free detergent fiber using the method described by⁴⁰.

Samples of okra fruits from each plot was taken for proximate analysis. The ash, crude protein, crude fat and carbohydrate contents of the okra fruits were determined using standard chemical methods described by Association of Analytical chemists³⁹. Samples of okra fruits were sliced with a knife and blended. After blending, it was diluted with ten times its weight with water (1:10). The viscous solution was separated from the debris using fine cloth⁴¹. Mucilage of the extracted viscous liquid was measured using viscometer.

About 2 g of each organic manures used was collected and analysed for N, P, K, Ca, and Mg as described by⁴². N was determined by the micro-Kjeldahl digestion method. Samples were digested with nitricperchloric-sulphuric acid mixture for the determination of P, K, Ca, and Mg. Phosphorus was determined colorimetrically using the vanadomolybdate method, K was determined using a flame photometer and Ca and Mg were determined by the EDTA titration method⁴³.

Statistical analysis. The data collected were subjected to statistical analysis of variance (ANOVA) using the Genstat statistical package⁴⁴ and treatment means were separated using Duncan Multiple Range Test (DMRT) at 5% probability level.

Received: 20 May 2020; Accepted: 27 August 2020

Published online: 30 September 2020

References

- Gemedo, H. F., Ratta, N., Haki, G. D., Woldegiorgis, A. Z. & Beyen, F. Nutritional quality and health benefits of “okra” (*Abelmoschus esculentus*): A review. *Int. J. Nutr. Food Sci.* **4**(2), 208–215 (2015).
- Adekiya, A. O., Aboyeji, C. M., Dunsin, O., Adebiyi, O. V. & Oyinlola, O. T. Effect of urea fertilizer and maize cob ash on soil chemical properties, growth, yield, and mineral composition of okra, *Abelmoschus esculentus* (L.) Moench. *J. Hortic. Res.* **26**(1), 67–76. <https://doi.org/10.2478/johr-2018-0008> (2018).
- Oyulu, C. Okra seeds: Potential source of high quality vegetable oil. in *Proceedings of 5th Annual Conference Horticulture Society, Nigeria*, Nsukka (1983).
- Adekiya, A. O., Agbede, T. M., Aboyeji, C. M., Dunsin, O. & Ugbe, J. O. Green manures and NPK fertilizer effects on soil properties, growth, yield, mineral and vitamin C composition of okra (*Abelmoschus esculentus*(L.) Moench). *J. Saudi Soc. Agric. Sci.* **18**, 218–223. <https://doi.org/10.1016/j.jssas.2017.05.005> (2019).
- Oladipo, O. G., Olayinka, A. & Aduayi, E. A. Effects of organic amendments on microbial activity, N and P mineralization in an Alfisol. *Environ. Manage. J.* **2**, 30–40 (2005).
- Maheswarappa, H. P., Nanjappa, H. V., Hegde, M. R. & Prabhu, S. R. Influence of planting material, plant population and organic manures on yield of East Indian galangal (*Kaempferia galanga*), soil physico-chemical and biological properties. *Indian J. Agron.* **44**(3), 651–657 (1999).
- Olowoake, A. A. Influence of organic, mineral and organomineral fertilizers on growth, yield, and soil properties in grain amaranth (*Amaranthus cruentus* L.). *J. Org.* **1**(1), 39–47 (2014).
- Agbede, T. M. & Adekiya, A. O. Effect of wood ash, poultry manure and NPK fertilizer on soil and leaf nutrient composition, growth and yield of okra (*Abelmoschus esculentus*). *Emirate J. Food Agric.* **24**(4), 314–321 (2012).
- Khandaker, M. M., Jusoh, N., Ralmi, N. H. A. & Ismail, S. Z. The effect of different types of organic fertilizers on growth and yield of *Abelmoschus esculentus* L. Moench (okra). *Bulg. J. Agric. Sci.* **23**(1), 119–125 (2017).
- Tiamiyu, R. A., Ahmed, H. G. & Muhammad, A. S. Effect of sources of organic manure on growth and yields of okra (*Abelmoschus esculentus* L.) in Sokoto, Nigeria. *Niger. J. Basic Appl. Sci.* **20**(3), 213–216 (2012).
- Fagwalawa, L. D. & Yahaya, S. M. Effect organic manure on the growth and yield of okra. *Imperial J. Interdiscipl. Res.* **2**(3), 130–133 (2016).
- Akinrinde, E. A. & Obigbesan, G. O. Evaluation of the fertility status of selected soils for crop production in five ecological zones of Nigeria. In *Proceedings of the 26th Annual Conference of Soil Science Society of Nigeria* (ed. Babalola, O.) 279–288 (University of Agriculture, Ibadan, 2000).
- Mbah, C. N. & Mbagwu, J. S. C. Effect of organic wastes on physicochemical properties up a dystrice leptosol and maize yield in southeastern—Nigeria. *Niger. J. Soil Sci.* **16**, 96–103 (2006).
- Adekiya, A. O. Legume mulch materials and poultry manure affect soil properties, and growth and fruit yield of tomato. *Agric. Consect. Sci.* **83**(2), 161–167 (2018).
- Agbede, T. M., Adekiya, A. O. & Ogeh, J. S. Response of soil properties and yam yield to *Chromolaena odorata* (Asteraceae) and *Tithonia diversifolia* (Asteraceae) mulches. *Arch. Agron. Soil Sci.* **60**(2), 209–224 (2014).
- Wolf, B. & Snyder, G. H. *Sustainable Soils: The Place of Organic Matter in Sustaining Soils and Their Productivity* (The Haworth Press Inc., New York, 2003).
- Togun, A. O., Akanbi, W. B. & Adediran, J. A. Growth, nutrient uptake and yield of tomato in response to different plant residue composts. *J. Food Agric. Environ.* **2**(1), 310–316 (2004).
- Olaniyi, J. O., Akanbi, W. B., Oladiran, O. A. & Ilupeju, O. T. The effect of organo-mineral and inorganic fertilizers on the growth, fruit yield, quality and chemical compositions of Okra. *J. Anim. Plant Sci.* **1**, 1135–1140 (2010).
- Ajari, O., Tsado, L. E. K., Oladiran, J. A. & Salako, E. A. Plant height and fruit yield of okra as affected by field application of fertilizer and organic matter in Bida, Nigeria. *Niger. Agric. J.* **34**, 74–80 (2003).
- Adekiya, A. O., Agbede, T. M., Aboyeji, C. M. & Dunsin, O. Response of okra (*Abelmoschus esculentus* (L.) Moench) and soil properties to different mulch materials in different cropping seasons. *Sci. Hortic.* **217**, 209–216 (2017).
- Adekiya, A. O., Agbede, T. M. & Ojaniyi, S. O. The effect of three years of tillage and poultry manure application on soil and plant nutrient composition, growth and yield of cocoyam. *Exp. Agric.* **52**, 466–476 (2016).
- Agbede, O. O. *Understanding Soil and Plant Nutrition* (Petra Digital Press, Abuja, 2009).
- Oyenuga, V. A. & Fetuga, B. I. Dietary importance of fruits and vegetables. in *Proceeding First National Seminar on Fruits and Vegetables*, 122–131. University of Ibadan. October 13–17 (1975).
- Rubatzky, V. E. & Yamaguchi, M. *World Vegetables: Principles, Production and Nutritive Values* 2nd edn. (International Thomas Publishing, Chapman and Hall, New York, 1997).
- Blumenthal, J., Battenspenrger, D., Cassman, K. G., Mason, K. G. & Pavlista, A. Importance of nitrogen on crop quality and health. In *Nitrogen in the Environment: Sources, Problems and Management* 2nd edn (eds Hatfield, J. L. & Folett, R. F.) (Elsevier, Amsterdam, 2008).
- Mani, S. & Ramanathan, K. M. Effect of nitrogen and potassium on the crude fibre content of bhendi fruit on successive stage of picking. *South Indian Hortic.* **29**(2), 100–104 (1981).
- Ahmad, E., Moaveni, P. & Farahani, H. A. Effects of planting dates and compost on mucilage variations in borage (*Borago officinalis* L.) under different chemical fertilization systems. *Int. J. Biotechnol. Mol. Biol. Res.* **1**(5), 58–61 (2010).
- Leroy, B. M. M., Bommele, L., Reheul, D., Moen, M. & de Neve, S. The application of vegetable, fruit and garden waste (VFG) compost in addition to cattle slurry in a silage maize monoculture: Effects on soil fauna and yield. *Eur. J. Soil Biol.* **43**, 91–100 (2007).
- Lumpkin, H. Organic vegetable production: A theme for international agricultural research. in *Proceedings of the seminar on the production and export of organic fruit and vegetables in Asia*. <https://www.fao.org/DOCREP/006/AD429E/ad429e13.htm>. Accessed 8 Dec 2019 (2003).
- Weston, L. A. & Barth, M. M. Pre-harvest factors affecting post-harvest quality of vegetables. *HortScience* **32**(5), 812–816 (1997).
- Lefsrud, M. G., Kopsell, D. A., Kopsell, D. E. & Curran-Celentano, J. Air temperature affect biomass and carotenoid pigment accumulation in kale and spinach grown in a controlled environment. *HortScience* **40**(7), 2026–2030 (2005).
- Cardoso, M. O. & Berni, R. F. Nitrogen applied in okra under non-tightness grown and residual fertilization. *Hortic. Bras.* **30**, 645–652 (2012).
- Gee, G. W. Particle-size analysis. In *Methods of Soil Analysis, Part 4. Physical Methods* (eds Dane, J. H. & Topp, G. C.) 255–293 (Wiley, Hoboken, 2002).

34. Nelson, D. W. & Sommers, L. E. Total carbon, organic carbon, and organic matter. In *Methods of Soil Analysis, Part 3. Chemical Methods* (ed. Sparks, D. L.) 961–1010 (Wiley, Hoboken, 1996).
35. Bremner, J. M. Nitrogen-total. In *Methods of Soil Analysis, Part 3. Chemical Methods* (ed. Sparks, D. L.) 1085–1121 (Wiley, Hoboken, 1996).
36. Frank, K., Beegle, D. & Denning, J. Phosphorus. In *Recommended chemical soil test procedures for the North Central Region, North Central Regional Research Publication No. 221 (revised)* (ed. Brown, J. R.) 21–26 (Missouri Agriculture Experiment Station, Columbia, 1998).
37. Hendershot, W. H. & Lalonde, H. Ion exchange and exchangeable cations. In *Soil Sampling and Methods of Analysis* (ed. Carter, M. R.) (Lewis Publishers, CRC Press, Cambridge, 1993).
38. Omolaiye, J. A. *et al.* Development of leaf area prediction model of okra (*Abelmoschus* spp.). *Product. Agric. Technol. J.* **11**(1), 130–136 (2015).
39. AOAC. *Official Methods of Analysis of AOAC International* (AOAC, Arlington, 2003).
40. Williams, P., El-Baramen, F. J., Nakkow, B. & Rihawi, S. *Crop Quality Evaluation Methods and Guidelines* (International Centre for Agricultural research in the Dry Area, Aleppo, 1986).
41. Thanatcha, R. & Pranee, A. Extraction and characterization of mucilage in *Ziziphus mauritiana* Lam. *Int. Food Res. J.* **18**, 201–212 (2011).
42. Tel, D.A. & Hagarty, M. *Soil and Plant Analysis. Study Guide for Agricultural Laboratory Directors and Technologists Working in Tropical Regions*. International Institute of Tropical Agriculture, Ibadan, Nigeria, in conjunction with the University of Guelph, Canada (1984).
43. Horwitz, W. & Latimer, G. W. (eds) *Official Methods of Analysis of AOAC International* (AOAC, Arlington, 2005).
44. Genstat. *Genstat 5 Release 3.2 Reference Manual* (Oxford University Press, Oxford, 1993).

Acknowledgements

I would like to thank Landmark University who provided enabling environment in carrying out the experiment and also willing to pay the article processing charges of this paper

Author contributions

A.O.A.—design, statistical analysis & write up; W.S.E.—typing of manuscript; A.O.—proof reading of manuscript; C.M.A.—design of experiment; O.D.—data collection; O.A.—field work; K.A.A.—proof reading of manuscript; C.A.—statistical analysis of data.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to A.O.A.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2020