



Prevalence and Antibiotic Resistance Profile of *Pseudomonas aeruginosa* from Hospital Sinks in South Western Nigeria

**Olayinka Oluyemi Oluranti¹, Chiamaka Ifeoma Ubanagu¹,
Olukunle Oluwapamilerin Oluwasemowo², Omowunmi Temidayo Akinola^{1,3},
Yewande Tolulope Nejo¹ and Babatunde Olanrewaju Motayo^{3*}**

¹Department of Biological Science, Bowen University Iwo, Nigeria.

²Department of Virology, University of Ibadan, Ibadan, Nigeria.

³Department of Pathology, Federal Medical Centre, Abeokuta, Nigeria.

Authors' contributions

This work was carried out in collaboration between all authors. Author OOO designed the study, authors CIU, OOO and BOM performed the statistical analysis. Author CIU and BOM wrote the protocol, and wrote the first draft of the manuscript. Authors OOO, CIU and OOO managed the analyses of the study. Authors CIU, YTN and OTA managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMMR/2019/v29i230067

Editor(s):

(1) Dr. Oyewole O. Olufemi, Department of Physiotherapy, Olabisi Onabanjo University Teaching Hospital, Nigeria.
(2) Dr. Muhammad Torequl Islam, Nuclear of Pharmaceutical Technology (NTF), Federal University of Piaui, Brazil.

Reviewers:

(1) Asitkumar Chakraborty, Vidyasagar University, India.

(2) Ronald Bartzatt, University of Nebraska, USA.

(3) Victor B. Oti, Nasarawa State University, Nigeria.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/47974>

Short Research Article

Received 22 December 2018

Accepted 15 March 2019

Published 25 March 2019

ABSTRACT

Background: *Pseudomonas aeruginosa* (*P. aeruginosa*) has been identified as a major pathogen in man, causing both opportunistic and nosocomial infections. *Pseudomonas* is a ubiquitous organism often isolated from various surfaces, which have the ability to form biofilms, making it a unique organism of medical importance.

Objective: The aim of this study is to determine the prevalence of *P. aeruginosa* isolated from hospital sinks and their antibiotic resistance profile.

Methods: Swab samples were collected from hospital sinks in five health care institutions and

*Corresponding author: E-mail: batatundemotayo@yahoo.com;

inoculated unto Nutrient agar and sub cultured on cetrimide agar. Isolated *P. aeruginosa* were subjected to antibiotic susceptibility testing using CSLI guidelines.

Results: Prevalence of *Pseudomonas* species isolated from the hospitals' sinks was 56%. High level resistance was recorded against amoxicillin/clavunilate, ampicillin and ceftriaxone. Resistance profile of the isolates clustered into two main clades clade A and clade B, with clade A isolates recording a higher MARI score.

Conclusion: Isolation of multi-resistant *P. aeruginosa* from hospital sinks calls for improved hospital infection control practices. We advocate for inclusion of environmental surveillance, particularly of opportunistic pathogens in our hospitals.

Keywords: Prevalence; *Pseudomonas aeruginosa*; antibiotic resistance; hospital sinks; MARI; Nigeria.

1. INTRODUCTION

Pseudomonas spp is a gram negative rod-like bacteria belonging to the family Pseudomonaceae, responsible for a host of infectious diseases in both man and animals such as Otitis externa, folliculitis, septicemia, and ventilator associated pneumonia [1]. *Pseudomonas* possesses certain properties such as quorum sensing and biofilm formation which allow it to spread and survive under adverse conditions [1]. Previous reports have also identified *Pseudomonas* spp to be responsible for outbreaks of dermatitis, conjunctivitis and otitis externa from recreational waters in the US [2]. Due to the ubiquitous nature of *Pseudomonas* spp it has been identified in various water sources [3,4,5], and as such water contributes to its cross contamination in clinical environments making it a primary agent of nosocomial infections [5,6].

Among organisms within *Pseudomonas* spp, *Pseudomonas aeruginosa* is the most medically important, it is responsible for chronic lung infections among cystic fibrosis patients [7]. *Pseudomonas aeruginosa* has also been identified as the singular most incriminated pathogen in nosocomial infection [8]. *Pseudomonas aeruginosa* is also unique for ability to acquire both plasmid borne and chromosomal antibiotic resistance genes [9]. Mutidrug resistance observed at high levels among *P. aeruginosa*, involves several mechanisms, including the overexpression of active efflux systems, production of beta-lactamase modifying enzymes, a decrease in outer membrane (OM) permeability as well as structural alterations of topoisomerases II and IV, involved in quinolone resistance [10,11]. In majority of MDR *P. aeruginosa*, some of these antibiotic-resistance genes are harbored within plasmids, clustered in a cassette carried by a

class 1 integron [11]. Fig. 1 represents an example of a gene map of an MDR plasmid showing the location of various antibiotic resistance genes. The *P. aeruginosa* genome, about 6.3 million bp encodes more than 8 virulence genes, making *P. aeruginosa* one of the most pathogenic bacteria with high mortality rates resulting from infections [12]. In Nigeria there have several reports on *P. aeruginosa* infections in clinical settings in Nigeria [8]. However there is a paucity of data on isolation of *P. spp* from fomites or contaminated surfaces such as sinks in hospital settings in Nigeria, despite its importance as a principal nosocomial pathogen able to survive on various surfaces and form biofilms. This study is aimed at determining the prevalence and antibiotic resistance profile of *P. aeruginosa* and other species of *Pseudomonas* in hospital sinks in some Nigerian hospitals.

2. MATERIALS AND METHODS

2.1 Study Design and Sample Collection

This study is a cross sectional survey of *Pseudomonas* species isolated from sinks in 5 tertiary health institutions in South western Nigeria, namely Lagos University Teaching Hospital (LUTH), University College Hospital, Ibadan (UCH), Nigeria Navy Reference Hospital, Lagos (NNRH), Pentecost Medical Centre, Lagos (PMC) and Bowen University Hospital, Iwo (BUH). Ethical approval was sought from the relevant ethics committee of the various facilities.

Sinks used for hand washing were randomly selected from the nursing department of each of the study facility, and swabbed with the aid of a sterile swab stick around the edges and around the tap knobs. Swab sticks were then transported aseptically to the Microbiology laboratory for sample processing.

2.2 Isolation of *Pseudomonas spp*

Samples were swabbed unto Nutrient/Mackonkay agar and incubated at 36°C for 48 hrs. Colonies were then subcultured on Cetrimide agar. Isolated pure cultures were then gram stained and tested using standard biochemical tests [12].

2.3 Antibiotic Susceptibility Testing

Antibiotic susceptibility testing was done on identified *Pseudomonas* isolates using commercially available single disks from Oxoid (U.K). Single colony of each isolate was inoculated into peptone water and 1.5 = 10 cfu/ml Macfarland suspension was inoculated unto Muller Hinton Agar. Single disks of Gentamicin (GEN), 10 µg; Ciprofloxacin (CPR), 5 µg; Ofloxacin (OFL), 5 µg; Amoxicillin/clavulanate (AUG), 30 µg; Nitrofurantoin (NIT), 300 µg; Ampicillin (AMP), 10 µg; Ceftazidime (CAZ), 30 µg and Cefuroxime (CRX), 30 µg. Agar plate were then incubated at 36°C for 24 hrs. Zone of inhibition were measured and interpreted following Clinical Laboratory Standards Institute guidelines [13].

2.4 Antibiotic Resistance Relatedness

Dendro UPGMA utility software was used to build a dendrogram, and the distance matrix used to calculate a similarity matrix and transform coefficients into distances using the Unweighted pair group method with arithmetic mean (UPGMA) algorithm [14].

2.5 Statistical Analysis

All the data generated was grouped into Tables, charts were drawn with the aid of Microsoft excel. analysis of data was done using Statistical Package for Social Sciences SPSS Vs 20.0.

3. RESULTS

The prevalence of *Pseudomonas* species isolated from the hospitals' sinks are shown in Table 1. From Lagos University Teaching Hospital, University College Hospital, Ibadan, Bowen University Hospital, Nigerian Navy Reference Hospital, Pentecost Medical Centre, the occurrence of *Pseudomonas* species were 85.71%, 57.14%, 57.14%, 50%, 20% respectively. The antibiotic sensitivity test is shown in Fig. 2 Most strains were sensitive to

gentamicin, ciprofloxacin and ofloxacin, while they showed relative resistance to other antibiotics including amoxicillin clavulanate, nitrofurantoin, ampicillin, ceftazidime and cefuroxime.

The antibiotic resistance profile of the isolates is shown in a dendrogram Fig. 3, indicating their clustering pattern and antibiotic multiresistant index (MARI). *Pseudomonas aeruginosa* recorded a MARI of between 0.25 to 0.625, while *Pseudomonas spp* recorded MARI of 0.025 to 1.0. The isolates also clustered into two main clades clade A and clade B, with clade A isolates recording a higher MARI score.

4. DISCUSSION

Pseudomonas remains one of the most important pathogens known to man, with its ubiquitous nature and ability to form biofilms [1], makes it an effective nosocomial pathogen. This study surveyed the prevalence and resistance profile of *Pseudomonas* isolated from hospital sinks in 5 different hospitals. The total prevalence of *Pseudomonas ssp* isolated was 56%, while LUTH recorded the highest number of isolations, while PMC recorded the lowest. This observation could be attributed to the size and number of consultations, LUTH for instance is one the largest teaching hospitals in Nigeria. Regardless of the level of infection control, some nosocomial pathogens such as pseudomonas can be introduced into the hospital environment accidentally by attending care givers. In our study, the highest level of antibiotic resistance was recorded in Cefuroxime, and Amoxicillin, with 88% and 82% respectively. This is a clear indication of high level beta lactamase resistance among our pseudomonas isolates. Previous studies have recorded a high level of resistance to 2rd generation chephalosporins [15,16]. Although there have been other reports of low resistance rates of 2nd generation chephalosporins to pseudomonas, such as Yetkin et al [17]. In our study there was also high level resistance observed against augumentin (amoxicillin/clavunalate) with 65% resistance. This high level resistance is indicative of the presence of extended spectrum beta lactamase (ESBL) among our isolates, even though ESBL testing was not conducted on the isolates; a resistance to both cephalosporin and augumentin shows the presence of ESBL enzyme [18-20]. *Pseudomonas aeruginosa*, resistance to b-lactam antibiotics and aminoglyco-sides has been reported to be associated with the




Fig. 2. Overall antibiotic sensitivity pattern of the isolated *Pseudomonas* species. The y-axis represents percentage susceptibility/resistance, x-axis represents antibiotics used for testing
 Antibiotic key: GEN-Gentamycin, CPR-Ciprofloxacin, OFL-Ofloxacin, AUG-Amoxicillin/Clavunalate, NIT-Nitrofurantoin, AMP-Ampicillin, CAZ-Ceftrazone, CRX-Ceftazidime




Fig. 3. Antibiotic resistant relatedness of *Pseudomonas* spp isolates obtained from hospital sinks with their respective multi-antibiotic resistance index (MARI)
 N.B: P.s represents *Pseudomonas* species, P.a represents *Pseudomonas aureginosa*

5. CONCLUSION

The current observation of high MARI is evidence of the wide spread dissemination of *Pseudomonas* spp harboring transmissible multi-resistant genes such as ESBL and Carbapenem resistance genes KPC in our environment. The detection of this pathogen in hospital sinks is worrisome, because *Pseudomonas* is able to form biofilm on various materials including plastic tubing used in most invasive medical devices such as catheter and naso-gastric tube. Effort should be made toward incorporating practices such as hospital environmental surveillance and effective environmental disinfection in various hospital infection prevention and control units.

ACKNOWLEDGEMENTS

The authors would like to acknowledge the management of the various hospitals where the study was carried out. We would also like to acknowledge Dr Paul Akinduti with his help in useful criticism and dendrogram construction.

CONSENT

It is not applicable.

ETHICAL APPROVAL

Ethical approval was sought from the relevant ethics committee of the various facilities.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Obritsch MD, Fish DN, McLaren R, Jung R. National surveillance of antimicrobial resistance in *Pseudomonas aeruginosa* isolates obtained from Intensive Care Unit Patients from 1993 to 2002. *Antimicrob Agents Chemother.* 2004;48(12):4606-4610.
2. Craun GF, Calderon RL, Craun MF. Outbreaks associated with recreational waters in the United States. *Int. J. Environ. Health. Res.* 2005;15 (4):243-62.
3. Ajayi AA, Shirdar MKC, Adekunle LV, Oluwande PA. Quality of packaged waters sold in Ibadan, Nigeria. *Afr. J. Biomed. Res.* 2008;11:251-258.
4. Barben J, Hafen G, Schmid J. *Pseudomonas aeruginosa* in public swimming pools and bathroom water of patients with cystic fibrosis. *Journal of Cystic Fibrosis.* 2005;4(4):227-231.
5. Hirulkar NB, Soni B. Incidence of antibiotic-resistant *Pseudomonas aeruginosa* isolated from drinking water. *Int. J. Pharm. Biol. Arch.* 2011.2(2):724-733.
6. Rueter S, Sigge A, Wiedect H, Trautmann M. Analysis of transmission pathways of *Pseudomonas aeruginosa* between patients and tap water outlets. *Critical Care Medicine.* 2002.30(10):2222-2228.
7. Pollack M. Principles and practice of infectious diseases, eds. In: Mandell GL, Bennet JE, Dollin R (Churchill Livingstone, Philadelphia). 2015;2:2310-2335.
8. Akingbade OA, Balogun SA, Ojo DA, Afolabi RO, Motayo BO, Okerentugba PO, Okonko IO. Plasmid profile analysis of multidrug resistant *Pseudomonas aeruginosa* isolated from wound infections in South West, Nigeria. *Wld. Appl. Sci. J.* 2012;20(6):766-775.
9. Adesoji AT, Ogunjobi AA, Olatoye IO. Molecular characterization of selected multidrug resistance *Pseudomonas* from water distribution systems in Southwestern Nigeria. *Ann Clinn Microbiol Antimicrob.* 2015;14(39):1-11.
10. Strateva T, Yordanov D. *Pseudomonas aeruginosa* a phenomenon of bacterial resistance. *J Med Microbiol.* 2009;58: 1133–1148.
11. Fuste E, Lopez-Jimenez L, Segura C, Gainza C. Carbapenem-resistance mechanisms of multidrug-resistance *Pseudomonas aeruginosa*. *J. Med. Microbiol.* 2013;62:1317-1325.
12. Agarwal G, Kapil A, Kabra SK, Das BK, Dwivedi N. Characterization of *Pseudomonas aeruginosa* isolated from chronically infected children with cystic fibrosis in India. *BMC Microbiology.* 2005; 5:43.
13. Cheesbrough M. District laboratory practice in tropical countries. Part 2. Cambridge University Press. 2006;434.
14. Clinical and Laboratory Standard Institute. Twenty-First Informational Supplement, CLSI Document M100-S21. Wayne, PA: Clinical and Laboratory Standards Institute. Performance Standards for Antimicrobial Susceptibility Testing; 2011.
15. Castillo-Vera J, Ribas-Aparicio RM, Nicolau CJ, Oliver A, Osorio-Carranza L,

- Aparicio-Ozores G. Unusual diversity of acquired β -lactamases in multidrug-resistant *Pseudomonas aeruginosa* isolates in a Mexican hospital. *Microbial Drug Resist.* 2012;18(5):471–478.
16. Olayinka AT, Olayinka BO, Onile BA. Antibiotic Susceptibility and Plasmid pattern of *Pseudomonas aeruginosa* from the surgical unit of a University Teaching Hospital in North Central, Nigeria. *Int. J. Med. Med. Sci.* 2009;1:79-83.
 17. Yetkin G, Otilu B, Cicek A, Kuzucu C, Durmaz R. Clinical microbiologic and epidemiologic characteristics of *P. aeruginosa* infections in a university hospital, Malaya, Turkey. *Am. J. Infect. Control.* 2006;34:188-192.
 18. Akinduti PA, Oluwadun A, Iwalokun B, Ejilude O, Onagbesan KO. Clonal dissemination of blaTEM beta-lactamase strains among enteric isolates in Abeokuta, Nigeria. *Res. J. Microbiol.* 2011;6(12):919-925.
 19. Olowe OA, Aboderin BW, Idris OO, Mabayoje VO, Opaleye OO, Adekunle OC, Olowe RA, Akinduti PA, Ojurongbe O. Genotypes of phenotypes of Shiga toxin-producing *Escherichia coli* (STEC) in Abeokuta, Southwestern Nigeria. *Infect. Drug. Resist.* 2014;7:253-259.
 20. Adeyakinu FA, Motayo BO, Akinduti PA, Akinbo J, Ogiogwa IJ, Aboderin BW, Agunlejika RA. A multicenter study of beta lactamase resistant *Escherichia coli* and *Klebsiella pneumoniae* reveals high level chromosome mediated beta-lactase resistance in Ogun State, Nigeria. *Interdisciplinary Perspectives on Infectious Diseases.* 2014;819896. DOI: 10.1155/2014/819896.
 21. Anupurba S, Bhattacharjee A, Garg A, Sen Malay R. Antimicrobial susceptibility of *Pseudomonas aeruginosa* isolated from wound infections. *Indian Journal of Dermatology.* 2006;51(4):286-288.
 22. Shittu OB, Adeniran SA, Afolabi OR, Sam-Wobo SO. Risk surveillance of multidrug resistant *Pseudomonas aeruginosa* in water and plasmid relatedness with clinical strains in Abeokuta, Southwest Nigeria. *J. Nat. Sci. Engr. & Tech.* 2014;13:44-57.
 23. Akinduti PA, Aboderin BW, Oloyede R, Ogiogwa JI, Motayo BO, Ejilude O. High-level multi-resistant and virulent *Escherichia coli* in Abeokuta, Nigeria. *J. Immunoassay. Immunochem.* 2016;37(2): 119-29.

© 2019 Oluranti et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/47974>